Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

SHOUT
SHARE
IT

UNITED STATES DEPARTMENT OF AGRICULTURE

DEPARTMENT BULLETIN No. 1352

Washington, D. C. August, 1925

# EFFECT OF AGE AND DEVELOPMENT ON BUTTERFAT PRODUCTION OF REGISTER-OF-MERIT JERSEY AND ADVANCED-REGISTER GUERNSEY CATTLE

By R. R. Graves, *Specialist in Dairy Cattle Breeding*, and M. H. Fohrman, *Dairy Husbandman, Bureau of Dairying*

CONTENTS

| | Page | | Page |
|---|---|---|---|
| The need for more information | 1 | Effect of pregnancy | 21 |
| Material used in this study | 2 | Recent improvement in production | 22 |
| Increase in production with age | 2 | Summary and conclusions | 22 |
| The factor of development | 5 | | |

## NEED FOR MORE INFORMATION

Age and development have a marked influence on the production of milk by dairy cows. This fact is acknowledged by the various record associations, and the requirements for recognition of cows as high producers are adjusted according to the age at which they are tested. The register of production is known in the Jersey breed as the Register of Merit, and in the Guernsey breed as the Advanced Register.

The student of the heredity of productive ability in dairy cattle who draws his material from the advanced-register and register-of-merit records is confronted with the problem of establishing some basis for comparing records made by cows of different ages, so as to avoid discrimination against the animal with a record made at an immature age. With the vast amount of official testing of dairy cattle that has been done there has accumulated a large fund of information which is useful and desirable for studies of this character. Cows and heifers are tested at various ages. Many animals tested as heifers are not retested after reaching maturity; consequently, without some method of adjusting records made at varying ages to a comparable basis, the use of this material is greatly restricted.

The value of a bull is largely based upon his ability to sire heifers of uniformly high production. In order to determine the transmitting ability of a sire, it is necessary to compare the records of his daughters with those of their dams. To limit the comparison only to those sets of dams and daughters with records at the same age

47588°—25†—Bull. 1352——1

would leave too few for positive conclusions, and to compare records of animals of different ages we must use some means of adjusting those of immature animals.

It is also a matter of interest to every breeder to know what relationship exists between the record of a 2-year animal and that of the same animal at 5 years of age, in order to be able to form some fairly accurate estimate of the ultimate ability of his heifers.

## MATERIAL USED FOR THIS STUDY

The sources of material for this study are the published results of official tests of Jersey and Guernsey cattle. The study embraces all class A and class AA records up to and including the 1921 volume of the Register of Merit of Jersey Cattle; and all single-letter advanced-register records up to and including Volume 34, No. 6 (April, 1922) of the Herd Register of the American Guernsey Cattle Club. A total of 14,571 Jersey records and 8,602 Guernsey records are used in the study of the effect of age on the production of butterfat.

These sources offer an abundance of authentic material for studies of this character. The records are made under careful supervision of the State experiment stations. The objections to using them are (1) that they are made by a selected group of individual animals, as the requirements for entry into the Advanced Register and Register of Merit exclude certain animals which fail to meet their requirements; (2) that cows on official test are usually those which the breeder thinks in advance will qualify; (3) and that before and during the test they receive better care and feed than ordinary herd cows.

It should be remembered, however, that a table of relationship of records of different ages evolved from this material would be used almost solely in connection with official records, because most studies of inheritance and prepotency of sires are based on records made under advanced-register conditions.

Though official records represent a selected group of animals, these animals were tested on hundreds of farms scattered across the country from coast to coast and from the Canadian line to the Gulf of Mexico. All conditions of climate are encountered in such a wide territory, systems of herd management range from everyday farm care to highly specialized treatment, and a great variety of feeds are used. The environmental effect of any particular locality or system of handling test cows is neutralized when all records are blended in an average drawn from so many animals scattered over a wide area.

## INCREASE IN PRODUCTION WITH AGE

The objects of this study were to determine (1) the increase in production that came with increasing age; (2) whether this increase in production with age was the same for younger as for older animals; and (3) at what age the maximum production was reached.

It was first planned to use only the records of animals that had two or more records, but the factor of development was found to enter into the second record in addition to the factor of increasing age, hence to determine the influence of age a study was made of initial records only. It may be assumed, therefore, that the difference between the average records for the various age groups shown in the tables is due to the influence of age only.

Tables 1 and 2 show what proportion the average production at any age is of the average mature production.

TABLE 1.—*Initial records at various ages of single-letter advanced-register Guernsey cows*

| Age (years) | Number of records | Average production of butterfat | Percentage that production at this age is of average mature production | Age (years) | Number of records | Average production of butterfat | Percentage that production at this age is of average mature production |
|---|---|---|---|---|---|---|---|
| | | *Pounds* | *Per cent* | | | *Pounds* | *Per cent* |
| Under 2 | 275 | 368 | 74.8 | 9 to 9½ | 102 | 502 | 102.0 |
| 2 to 2½ | 2,367 | 404 | 82.1 | 9½ to 10 | 80 | 496 | 100.8 |
| 2½ to 3 | 1,211 | 424 | 86.2 | 10 to 10½ | 45 | 493 | 100.2 |
| 3 to 3½ | 897 | 433 | 88.0 | 10½ to 11 | 27 | 486 | 98.8 |
| 3½ to 4 | 695 | 453 | 92.1 | 11 to 11½ | 28 | 459 | 93.3 |
| 4 to 4½ | 625 | 470 | 95.5 | 11½ to 12 | 15 | 456 | 92.7 |
| 4½ to 5 | 432 | 473 | 96.1 | 12 to 12½ | 13 | 504 | 102.4 |
| 5 to 5½ | 373 | 495 | 100.6 | 12½ to 13 | 8 | 475 | 96.5 |
| 5½ to 6 | 313 | 489 | 99.4 | 13 to 13½ | 10 | 454 | 92.3 |
| 6 to 6½ | 305 | 482 | 98.0 | 13½ to 14 | 8 | 494 | 100.4 |
| 6½ to 7 | 214 | 487 | 99.0 | 14 to 14½ | 3 | 417 | 84.8 |
| 7 to 7½ | 179 | 495 | 100.6 | 15 to 15½ | 3 | 456 | 92.7 |
| 7½ to 8 | 148 | 499 | 101.4 | 16½ to 17 | 1 | 395 | 80.3 |
| 8 to 8½ | 135 | 496 | 100.8 | 17 to 17½ | 1 | 458 | 93.1 |
| 8½ to-9 | 88 | 495 | 100.6 | 17½ to 18 | 1 | 364 | 74.0 |

TABLE 2.—*Initial records at various ages of class A and class AA register-of-merit Jersey cows*

| Age (years) | Number of records | Average production of butterfat | Percentage that production at this age is of average mature production | Age (years) | Number of records | Average production of butterfat | Percentage that production at this age is of average mature production |
|---|---|---|---|---|---|---|---|
| | | *Pounds* | *Per cent* | | | *Pounds* | *Per cent* |
| Under 2 | 1,237 | 343 | 72.7 | 9½ to 10 | 128 | 475 | 100.6 |
| 2 to 2½ | 3,633 | 368 | 78.0 | 10 to 10½ | 114 | 476 | 100.8 |
| 2½ to 3 | 1,313 | 386 | 81.8 | 10½ to 11 | 68 | 459 | 97.2 |
| 3 to 3½ | 1,298 | 404 | 85.6 | 11 to 11½ | 53 | 454 | 96.2 |
| 3½ to 4 | 1,011 | 423 | 89.8 | 11½ to 12 | 39 | 478 | 101.3 |
| 4 to 4½ | 924 | 440 | 93.2 | 12 to 12½ | 40 | 445 | 94.3 |
| 4½ to 5 | 705 | 454 | 96.2 | 12½ to 13 | 16 | 436 | 92.4 |
| 5 to 5½ | 673 | 457 | 96.8 | 13 to 13½ | 18 | 445 | 94.3 |
| 5½ to 6 | 587 | 469 | 99.4 | 13½ to 14 | 11 | 475 | 100.6 |
| 6 to 6½ | 509 | 475 | 100.6 | 14 to 14½ | 6 | 439 | 93.0 |
| 6½ to 7 | 404 | 471 | 99.8 | 14½ to 15 | 6 | 424 | 89.8 |
| 7 to 7½ | 395 | 468 | 99.2 | 15 to 15½ | 2 | 412 | 87.3 |
| 7½ to 8 | 310 | 470 | 99.6 | 15½ to 16 | 4 | 405 | 85.8 |
| 8 to 8½ | 267 | 480 | 101.7 | 16 to 16½ | 2 | 437 | 92.6 |
| 8½ to 9 | 232 | 462 | 97.9 | 16½ to 17 | 2 | 391 | 82.8 |
| 9 to 9½ | 163 | 468 | 99.2 | 18 to 18½ | 1 | 401 | 85.0 |

The data studied do not show that there is a definite age at which all cows reach their maximum producing ability. For both Guernsey and Jersey cattle there is a period of years during which the average production of animals in the various age groups shows only slight variation. These fluctuations have no decided trend, and this entire period is included in the age of maximum production. The Jersey cows seem to reach maturity at 6 years of age, and there is no decided decline in average production until after the age of 10½ years. There are 2,522 initial records in this period, according to Table 2,

and the average is 472 pounds of butterfat. The Guernsey cows seem to reach mature producing ability at 5 years of age, and there is no marked decline in the average production until the age of 11 years. As shown in Table 1, this 6-year period includes 2,009 initial records averaging 492 pounds of butterfat. The percentages shown in Tables 1 and 2 are based on these amounts as average mature production.

Unless this age of mature or maximum producing capacity is spread over a period of years it is difficult to account for the performances of those cows which have completed two or more records of approximately equal quantities of butterfat during mature life.

The distribution of the initial records for the two breeds studied is shown in Tables 3 and 4. It is true, however, that the requirements for entry of cows in the Register of Merit and Advanced Register exert a selective influence on the distribution of records by excluding animals which produce less than the required quantity of butterfat; and in addition there is some selection by breeders because of the practice of testing only certain animals in their herds.

The distribution is not symmetrical; 56.4 per cent of all the Guernsey records and 58 per cent of all the Jersey records fall below the averages of their respective groups.

On statistical analysis the 14,571 Jersey records show a mean of $411.41 \pm 0.51$ pounds of butterfat, with a standard deviation of $90.68 \pm 0.36$ pounds of fat, and a coefficient of variability of $22.04 \pm 0.09$.

The mean of the 8,602 Guernsey records was found to be $442.07 \pm 0.69$ pounds of butterfat, standard deviation $94.54 \pm 0.49$ pounds of butterfat, and coefficient of variability $21.39 \pm 0.11$.

TABLE 3.—*Distribution of initial records of single-letter advanced-register Guernsey cows according to age*

| Pounds of butterfat | Age | | | | | | | | | | Total |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | Under 2 years | 2 years | 3 years | 4 years | 5 years | 6 years | 7 years | 8 years | 9 years | 10 years and older | |
| 250 to 274 | 19 | 60 | | | | | | | | | 79 |
| 275 to 299 | 29 | 187 | 7 | | | | | | | | 223 |
| 300 to 324 | 44 | 325 | 85 | 1 | | | | | | | 455 |
| 325 to 349 | 35 | 383 | 142 | 34 | | | | | | | 594 |
| 350 to 374 | 31 | 436 | 170 | 95 | 34 | 30 | 13 | 13 | 9 | 14 | 845 |
| 375 to 399 | 31 | 415 | 193 | 122 | 65 | 55 | 23 | 26 | 14 | 19 | 963 |
| 400 to 424 | 30 | 390 | 179 | 114 | 77 | 60 | 29 | 16 | 20 | 19 | 934 |
| 425 to 449 | 16 | 336 | 145 | 112 | 72 | 66 | 46 | 30 | 23 | 21 | 867 |
| 450 to 474 | 17 | 271 | 165 | 129 | 76 | 65 | 34 | 20 | 26 | 21 | 824 |
| 475 to 499 | 10 | 210 | 120 | 106 | 85 | 61 | 36 | 19 | 16 | 6 | 669 |
| 500 to 524 | 7 | 181 | 93 | 88 | 62 | 42 | 36 | 25 | 16 | 16 | 566 |
| 525 to 549 | 3 | 135 | 105 | 62 | 43 | 28 | 35 | 13 | 13 | 14 | 451 |
| 550 to 574 | 1 | 91 | 62 | 61 | 49 | 29 | 22 | 17 | 6 | 14 | 352 |
| 575 to 599 | 2 | 52 | 44 | 39 | 35 | 28 | 17 | 15 | 12 | 5 | 249 |
| 600 to 624 | | 40 | 28 | 31 | 30 | 19 | 11 | 10 | 4 | 6 | 179 |
| 625 to 649 | | 24 | 25 | 23 | 25 | 12 | 8 | 4 | 8 | 3 | 132 |
| 650 to 674 | | 17 | 10 | 8 | 7 | 6 | 4 | 4 | 6 | | 62 |
| 675 to 699 | | 6 | 8 | 9 | 11 | 4 | 5 | 4 | 2 | 3 | 52 |
| 700 to 724 | | 8 | 4 | 5 | 10 | 9 | 3 | 3 | | 1 | 43 |
| 725 to 749 | | 7 | 3 | 5 | 2 | 1 | 1 | 2 | 1 | | 22 |
| 750 to 774 | | 2 | 2 | 5 | 2 | | | | 1 | 1 | 13 |
| 775 to 799 | | 1 | 1 | 4 | | 1 | 2 | 1 | 2 | | 12 |
| 800 to 824 | | 1 | | 3 | | 1 | 2 | | | | 7 |
| 825 to 849 | | | | 1 | 1 | 1 | | | | | 3 |
| 850 to 874 | | | 1 | | | | | | 2 | | 3 |
| 875 to 899 | | | | | | | | 1 | 1 | | 2 |
| 900 to 924 | | | | | | | | | | | |
| 925 to 949 | | | | | | 1 | | | | | 1 |
| Total | 275 | 3,578 | 1,592 | 1,057 | 686 | 519 | 327 | 223 | 182 | 163 | 8,602 |

TABLE 4.—*Distribution of initial records of class A and class AA Jersey cows, according to age*

| Pounds of butterfat | Age | | | | | | | | | | Total |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | Under 2 years | 2 years | 3 years | 4 years | 5 years | 6 years | 7 years | 8 years | 9 years | 10 years and older | |
| 250 to 274 | 172 | 231 | | | | | | | | | 403 |
| 275 to 299 | 183 | 611 | 29 | | | | | | | | 823 |
| 300 to 324 | 226 | 744 | 211 | 1 | | | | | | | 1,182 |
| 325 to 349 | 172 | 726 | 327 | 94 | | | | | | | 1,319 |
| 350 to 374 | 162 | 734 | 344 | 213 | 129 | 78 | 54 | 31 | 19 | 53 | 1,817 |
| 375 to 399 | 119 | 618 | 340 | 233 | 173 | 129 | 101 | 62 | 41 | 49 | 1,865 |
| 400 to 424 | 67 | 454 | 311 | 215 | 187 | 136 | 117 | 80 | 48 | 59 | 1,674 |
| 425 to 449 | 53 | 328 | 224 | 198 | 176 | 111 | 97 | 62 | 44 | 49 | 1,342 |
| 450 to 474 | 31 | 244 | 192 | 185 | 146 | 84 | 66 | 61 | 33 | 38 | 1,080 |
| 475 to 499 | 12 | 173 | 114 | 139 | 121 | 76 | 64 | 56 | 23 | 42 | 820 |
| 500 to 524 | 15 | 125 | 97 | 109 | 84 | 78 | 57 | 37 | 15 | 26 | 643 |
| 525 to 549 | 9 | 90 | 63 | 64 | 67 | 46 | 32 | 35 | 22 | 14 | 442 |
| 550 to 574 | 6 | 53 | 41 | 63 | 45 | 48 | 29 | 23 | 9 | 10 | 327 |
| 575 to 599 | 4 | 43 | 33 | 36 | 40 | 33 | 24 | 12 | 9 | 11 | 245 |
| 600 to 624 | 1 | 26 | 25 | 32 | 32 | 31 | 21 | 7 | 9 | 5 | 189 |
| 625 to 649 | 2 | 15 | 20 | 17 | 16 | 20 | 8 | 13 | 5 | 11 | 127 |
| 650 to 674 | 1 | 13 | 16 | 9 | 12 | 11 | 9 | 7 | 2 | 5 | 85 |
| 675 to 699 | | 9 | 8 | 4 | 10 | 8 | 5 | 1 | 3 | 2 | 50 |
| 700 to 724 | | 4 | 5 | 9 | 10 | 10 | 9 | 5 | 4 | 4 | 60 |
| 725 to 749 | | 1 | 4 | 3 | 3 | 5 | 3 | 3 | 2 | 1 | 25 |
| 750 to 774 | | 3 | 3 | 2 | 5 | 1 | 1 | 1 | | | 16 |
| 775 to 799 | | | | | 1 | 5 | 3 | 1 | | 1 | 11 |
| 800 to 824 | 1 | 1 | | 1 | 2 | 1 | 4 | | 2 | | 12 |
| 825 to 849 | 1 | | 1 | 2 | | 1 | | 2 | | | 7 |
| 850 to 874 | | | | | | | 1 | | | 1 | 2 |
| 875 to 899 | | | | | | | | | | | |
| 900 to 924 | | | | | | | | | | 1 | 1 |
| 925 to 949 | | | | | | 1 | | | | | 1 |
| 950 to 974 | | | | | 1 | | | | 1 | | 2 |
| 975 to 999 | | | 1 | | | | | | | | 1 |
| Total | 1,237 | 5,246 | 2,409 | 1,629 | 1,260 | 913 | 705 | 499 | 291 | 382 | 14,571 |

## THE FACTOR OF DEVELOPMENT

In studying the official-test records of animals grouped according to age it is apparent that in advancing from class to class there is some influence at work, other than age, which causes an increase in production. A comparison of the tables of initial records with those made up of both initial and reentry records brings this point out clearly. It is also brought to our attention in comparing records made at advanced ages by the same individuals, after age itself has ceased to bring about a further increase in production. With very few exceptions cows which made their initial records after reaching maturity have increased these records when tested again.

The first grouping of animals to study the influence of age on production was made only from animals having two or more records. It was found that because of the influence on the second record of the development made by the animal during the initial test, these retest records did not offer a correct indication of the effect on production of advance in age; but, when compared with the average initial records made for each age class, these retest records do show the apparent influence on later records of the development made by the animal during the first test.

Tables 5 to 21, inclusive, show the average initial and retest records of all Guernsey and Jersey cows having two records. A study of these tables will show the manner in which the animals are grouped. In Table 5, for instance, all the cows made their initial records under

2 years of age, and two cows in this class, constituting the first group, whose initial records average 295 pounds of butterfat, were retested when 2½ to 3 years old and made retest records averaging 393 pounds of butterfat. Fourteen cows with initial records averaging 396 pounds of butterfat were retested when 3 to 3½ years old and made retest records averaging 511 pounds of butterfat.

The difference between the average reentry and average initial records is divided into two parts, one of which represents the increment due to age and the other the portion of increase due to the development brought about by reason of the initial record.

The column which succeeds the series of reentry record columns shows the portion of the reentry record which is due to the increase in age since the initial record was made. The next double column in the table shows the portion of the reentry record which is due to development. This is given in pounds of butterfat and percentage of the reentry record. The last column shows the average per cent of increase due to development for the entire group.

To complete the illustration given above, the average reentry record of the two cows in the first group was 98 pounds greater than their initial record; 45 pounds of this increase was due to the year's increase in age, and 53 pounds, or 13.5 per cent, was due to the development the heifers underwent because of the feeding and handling during the previous test.

The first group in Table 13 includes 51 heifers with average initial records, made under 2 years of age, of 340 pounds of butterfat. These same 51 heifers made reentry records at 2½ to 3 years of age which average 412 pounds of butterfat. The total increase is 72 pounds of butterfat, of which 43 pounds is due to the year's increase in age, and 28 pounds, or 7 per cent, is due to development.

A word of explanation should be given as to how these results are obtained. The 51 Jerseys under 2 years of age in the first group of Table 13 average 340 pounds of butterfat on their initial tests. The average initial record of 1,237 Jerseys as shown in Table 2 is 343 pounds of butterfat. Therefore this group of 51 produced 99.1 per cent of the average of all animals under 2 years. The average production of 1,613 Jerseys at the age of 2½ to 3 years is 386 pounds of butterfat (Table 2), and 99.1 per cent of 386 is 383 pounds of butterfat, which should be the estimated production of this group when 2½ to 3 years old, allowing for the normal increase due to age. The difference between 412 pounds, the actual reentry record, and 383 pounds, the estimated production for this group at 2½ to 3 years of age, is 28 pounds, and this may be said to represent that portion of the increase which is due to development.

In this study there are 209 groups of Jerseys, and all but 23 of the groups show a positive development factor. There are 3,722 individuals in these 209 groups, and the 23 negative groups number only 39 individuals. These negative results are frequently based on groups numbering only one individual.

The Guernseys with two records number 1,215, listed in 148 groups. Only 14 individuals in 9 groups fail to show a positive development factor.

In a total of 3,722 pairs of Jersey records studied, the development factor averages 11.0 per cent of the reentry record; and in 1,215 pairs of Guernsey records an average of 12.2 per cent of the reentry record is due to development.

Where the groups are sufficiently large to be significant, this factor of development is fairly uniform. Groups containing small numbers of individuals show the greatest fluctuations, which is to be expected. The tables do seem to indicate, however, that where the interval is very long between the initial and reentry records the development is more marked. This may be contrary to what would be expected, or it is possible that during the lapse of a considerable period of time there has been sufficiently marked improvement in methods of handling test cows to account for this greater development.

There is sufficient evidence to prove that official testing develops the productive ability of dairy cows, and that the feeding and care, combined with the prolonged milking period during which the cow is encouraged to yield her utmost, have an appreciable and positive effect on her production during subsequent lactation periods.

TABLE 5.—*Comparison of average initial and retest records of Guernseys having initial records made at age under 2 years (single-letter records)*

| Number of cows | Average initial butterfat record made under 2 years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 2½ to 3 | 3 to 3½ | 3½ to 4 | 4 to 4½ | 4½ to 5 | 5 to 5½ | 5½ to 6 | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | | | | |
| | *Pounds* | | | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *P. ct.* |
| 2 | 295 | 393 | | | | | | | | | | | | | | | | 45 | 53 | 13. 5 | |
| 14 | 396 | | 511 | | | | | | | | | | | | | | | 70 | 45 | 8. 8 | |
| 4 | 358 | | | 463 | | | | | | | | | | | | | | 83 | 22 | 4. 8 | |
| 4 | 371 | | | | 538 | | | | | | | | | | | | | 103 | 64 | 11. 9 | |
| 6 | 388 | | | | | 578 | | | | | | | | | | | | 111 | 79 | 13. 7 | |
| 1 | 584 | | | | | | 695 | | | | | | | | | | | 197 | −86 | −12. 4 | |
| 7 | 383 | | | | | | | 559 | | | | | | | | | | 129 | 47 | 8. 4 | 8. 11 |
| 2 | 321 | | | | | | | | 467 | | | | | | | | | 108 | 38 | 8. 1 | |
| 4 | 407 | | | | | | | | | 571 | | | | | | | | 137 | 27 | 4. 7 | |
| 5 | 392 | | | | | | | | | | 486 | | | | | | | 132 | −38 | −7. 8 | |
| 1 | 321 | | | | | | | | | | | 533 | | | | | | 108 | 104 | 19. 5 | |
| 2 | 415 | | | | | | | | | | | | 603 | | | | | 140 | 48 | 8. 0 | |
| 2 | 311 | | | | | | | | | | | | | 643 | | | | 105 | 227 | 35. 3 | |
| 1 | 321 | | | | | | | | | | | | | | | | 422 | 108 | −7 | −1. 7 | |

TABLE 6.—*Comparison of average initial and retest records of Guernseys having initial records made at age of 2 years (single-letter records)*

| Number of cows | Average initial butterfat record made at 2 to 2½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 3 to 3½ | 3½ to 4 | 4 to 4½ | 4½ to 5 | 5 to 5½ | 5½ to 6 | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | | | |
| | *Pounds* | | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *P. ct.* |
| 46 | 424 | 501 | | | | | | | | | | | | | | | 31 | 46 | 9. 2 | |
| 106 | 449 | | 546 | | | | | | | | | | | | | | 54 | 43 | 7. 9 | |
| 50 | 427 | | | 541 | | | | | | | | | | | | | 70 | 44 | 8. 1 | |
| 70 | 435 | | | | 552 | | | | | | | | | | | | 74 | 43 | 7. 8 | |
| 36 | 418 | | | | | 596 | | | | | | | | | | | 91 | 87 | 14. 6 | |
| 34 | 416 | | | | | | 574 | | | | | | | | | | 91 | 67 | 11. 7 | |
| 34 | 430 | | | | | | | 600 | | | | | | | | | 93 | 77 | 12. 8 | |
| 22 | 381 | | | | | | | | 574 | | | | | | | | 83 | 110 | 19. 2 | 10. 7 |
| 18 | 406 | | | | | | | | | 587 | | | | | | | 88 | 93 | 15. 8 | |
| 11 | 443 | | | | | | | | | | 620 | | | | | | 97 | 80 | 12. 9 | |
| 6 | 401 | | | | | | | | | | | 576 | | | | | 88 | 87 | 15. 1 | |
| 10 | 402 | | | | | | | | | | | | 595 | | | | 88 | 105 | 17. 6 | |
| 2 | 390 | | | | | | | | | | | | | 637 | | | 85 | 162 | 25. 4 | |
| 2 | 333 | | | | | | | | | | | | | | 565 | | 72 | 160 | 28. 3 | |
| 2 | 390 | | | | | | | | | | | | | | | 541 | 85 | 66 | 12. 2 | |
| | Made at 2½ to 3 years old | | | | | | | | | | | | | | | | | | | |
| 43 | 448 | | 529 | | | | | | | | | | | | | | 31 | 50 | 9. 5 | |
| 47 | 447 | | | 544 | | | | | | | | | | | | | 48 | 49 | 9. 0 | |
| 27 | 434 | | | | 547 | | | | | | | | | | | | 50 | 63 | 11. 5 | |
| 23 | 455 | | | | | 558 | | | | | | | | | | | 73 | 30 | 5. 4 | |
| 19 | 447 | | | | | | 605 | | | | | | | | | | 72 | 86 | 14. 2 | |
| 17 | 423 | | | | | | | 573 | | | | | | | | | 68 | 82 | 14. 3 | |
| 16 | 409 | | | | | | | | 535 | | | | | | | | 66 | 60 | 11. 2 | 10. 6 |
| 8 | 409 | | | | | | | | | 579 | | | | | | | 66 | 104 | 18. 0 | |
| 4 | 423 | | | | | | | | | | 579 | | | | | | 68 | 88 | 15. 2 | |
| 6 | 489 | | | | | | | | | | | 578 | | | | | 78 | 11 | 1. 9 | |
| 2 | 359 | | | | | | | | | | | | 621 | | | | 58 | 204 | 32. 9 | |
| 2 | 496 | | | | | | | | | | | | | 647 | | | 80 | 71 | 11. 0 | |
| 1 | 333 | | | | | | | | | | | | | | | 519 | 53 | 133 | 25. 6 | |

TABLE 7.—*Comparison of average initial and retest records of Guernseys having initial records made at age of 3 years (single-letter records)*

| Number of cows | Average initial butterfat record made at 3 to 3½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 4 to 4½ | 4½ to 5 | 5 to 5½ | 5½ to 6 | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | | | |
| | *Pounds* | | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *Per cent* |
| 20 | 421 | 550 | | | | | | | | | | | | | | | 36 | 93 | 16. 9 | |
| 30 | 442 | | 567 | | | | | | | | | | | | | | 41 | 84 | 14. 8 | |
| 18 | 422 | | | 574 | | | | | | | | | | | | | 58 | 94 | 16. 4 | |
| 24 | 459 | | | | 565 | | | | | | | | | | | | 63 | 43 | 7. 6 | |
| 14 | 462 | | | | | 572 | | | | | | | | | | | 63 | 47 | 8. 2 | |
| 13 | 442 | | | | | | 563 | | | | | | | | | | 60 | 61 | 10. 8 | |
| 12 | 433 | | | | | | | 590 | | | | | | | | | 59 | 98 | 16. 6 | |
| 4 | 357 | | | | | | | | 474 | | | | | | | | 48 | 69 | 14. 6 | 13. 2 |
| 3 | 592 | | | | | | | | | 692 | | | | | | | 81 | 19 | 2. 7 | |
| 3 | 421 | | | | | | | | | | 637 | | | | | | 57 | 159 | 25. 0 | |
| 2 | 486 | | | | | | | | | | | 625 | | | | | 66 | 73 | 11. 7 | |
| 1 | 405 | | | | | | | | | | | | 774 | | | | 55 | 314 | 40. 6 | |
| 1 | 555 | | | | | | | | | | | | | 545 | | | 76 | —86 | —15. 8 | |
| 1 | 517 | | | | | | | | | | | | | | | 673 | 42 | 114 | 16. 9 | |
| | Made at 3½ to 4 years old | | | | | | | | | | | | | | | | | | | |
| 17 | 479 | | 556 | | | | | | | | | | | | | | 21 | 56 | 10. 0 | |
| 19 | 443 | | | 546 | | | | | | | | | | | | | 38 | 65 | 11. 9 | |
| 10 | 439 | | | | 537 | | | | | | | | | | | | 38 | 60 | 11. 2 | |
| 15 | 496 | | | | | 634 | | | | | | | | | | | 43 | 95 | 15. 0 | |
| 8 | 504 | | | | | | 601 | | | | | | | | | | 44 | 53 | 8. 8 | |
| 12 | 456 | | | | | | | 564 | | | | | | | | | 39 | 69 | 12. 2 | |
| 5 | 479 | | | | | | | | 705 | | | | | | | | 41 | 185 | 26. 2 | 14. 6 |
| 4 | 427 | | | | | | | | | 561 | | | | | | | 37 | 97 | 17. 3 | |
| 3 | 426 | | | | | | | | | | 611 | | | | | | 36 | 149 | 24. 4 | |
| 2 | 458 | | | | | | | | | | | 679 | | | | | 39 | 182 | 26. 8 | |
| 2 | 428 | | | | | | | | | | | | 683 | | | | 37 | 218 | 31. 9 | |
| 1 | 368 | | | | | | | | | | | | | 721 | | | 32 | 321 | 44. 5 | |
| 1 | 373 | | | | | | | | | | | | | | | 480 | 12 | 95 | 19. 8 | |

TABLE 8.—*Comparison of average initial and retest records of Guernseys having initial records made at age of 4 years (single-letter records)*

| Number of cows | Average initial butterfat record made at 4 to 4½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 5 to 5½ | 5½ to 6 | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | 11½ to 12 | | | | |
| | *Pounds* | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *Per cent* | *Per cent* |
| 13 | 501 | 607 | | | | | | | | | | | | | | 23 | 83 | 13.7 | |
| 14 | 506 | | 580 | | | | | | | | | | | | | 24 | 50 | 8.6 | |
| 10 | 457 | | | 626 | | | | | | | | | | | | 21 | 148 | 23.6 | |
| 16 | 464 | | | | 566 | | | | | | | | | | | 22 | 80 | 14.1 | |
| 8 | 442 | | | | | 516 | | | | | | | | | | 20 | 54 | 10.5 | |
| 1 | 401 | | | | | | 425 | | | | | | | | | 19 | 5 | 1.2 | |
| 4 | 473 | | | | | | | 574 | | | | | | | | 22 | 79 | 13.8 | 13.3 |
| 2 | 471 | | | | | | | | 575 | | | | | | | 22 | 82 | 14.3 | |
| 3 | 481 | | | | | | | | | 539 | | | | | | 22 | 36 | 6.7 | |
| 2 | 394 | | | | | | | | | | 484 | | | | | 18 | 72 | 14.9 | |
| 1 | 507 | | | | | | | | | | | 590 | | | | 24 | 59 | 10.0 | |
| 1 | 507 | | | | | | | | | | | | | 477 | | −2 | −28 | −5.9 | |
| 1 | 346 | | | | | | | | | | | | | | 481 | −2 | 137 | 28.5 | |
| | Made at 4½ to 5 years old | | | | | | | | | | | | | | | | | | |
| 9 | 442 | | 540 | | | | | | | | | | | | | 18 | 80 | 14.8 | |
| 9 | 526 | | | 606 | | | | | | | | | | | | 21 | 59 | 9.7 | |
| 4 | 517 | | | | 725 | | | | | | | | | | | 21 | 187 | 25.8 | |
| 8 | 559 | | | | | 683 | | | | | | | | | | 23 | 101 | 14.8 | |
| 3 | 506 | | | | | | 636 | | | | | | | | | 20 | 110 | 17.3 | |
| 4 | 452 | | | | | | | 593 | | | | | | | | 18 | 123 | 20.7 | 16.0 |
| 2 | 420 | | | | | | | | 561 | | | | | | | 17 | 124 | 22.1 | |
| 1 | 386 | | | | | | | | | 594 | | | | | | 15 | 193 | 32.5 | |
| 1 | 445 | | | | | | | | | | | | 462 | | | 18 | −1 | −0.2 | |
| 1 | 483 | | | | | | | | | | | | | | 601 | −5 | 123 | 20.5 | |

TABLE 9.—*Comparison of average initial and retest records of Guernseys having initial records made at age of 5 years (single-letter records)*

| Number of cows | Average initial butterfat record made at 5 to 5½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | | | | |
| | *Pounds* | | | | | | | | | *Pounds* | *Pounds* | *Per cent* | *Per cent* |
| 7 | 534 | 594 | | | | | | | | 0 | 60 | 10.1 | |
| 7 | 517 | | 541 | | | | | | | 0 | 24 | 4.4 | |
| 2 | 496 | | | 498 | | | | | | 0 | 2 | 0.4 | |
| 1 | 552 | | | | 571 | | | | | 0 | 19 | 3.3 | 9.2 |
| 3 | 458 | | | | | 579 | | | | 0 | 121 | 20.9 | |
| 1 | 404 | | | | | | 487 | | | 0 | 83 | 17.0 | |
| 1 | 454 | | | | | | | 555 | | 0 | 101 | 18.1 | |
| | Made at 5½ to 6 years old | | | | | | | | | | | | |
| 5 | 500 | | 546 | | | | | | | 0 | 46 | 8.4 | |
| 4 | 549 | | | 644 | | | | | | 0 | 95 | 14.8 | |
| 1 | 561 | | | | 710 | | | | | 0 | 149 | 21.0 | |
| 8 | 494 | | | | | 592 | | | | 0 | 98 | 16.6 | 16.8 |
| 2 | 477 | | | | | | 683 | | | 0 | 206 | 30.2 | |
| 1 | 438 | | | | | | | 657 | | 0 | 219 | 33.3 | |
| 2 | 492 | | | | | | | | 608 | 0 | 116 | 19.1 | |

TABLE 10.—*Comparison of average initial and retest records of Guernseys having initial records made at age of 6 years (single-letter records)*

| Number of cows | Average initial butterfat record made at 6 to 6½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | 11½ to 12 | | | | |
| | *Pounds* | | | | | | | | | | | *Pounds* | *Pounds* | *Per cent* | *Per cent* |
| 5 | 500 | 723 | | | | | | | | | | 0 | 223 | 30.8 | |
| 6 | 468 | | 607 | | | | | | | | | 0 | 139 | 22.9 | |
| 2 | 419 | | | 417 | | | | | | | | 0 | −2 | −0.5 | |
| 3 | 567 | | | | 701 | | | | | | | 0 | 134 | 19.1 | |
| 4 | 475 | | | | | 677 | | | | | | 0 | 202 | 29.8 | |
| 2 | 411 | | | | | | 543 | | | | | 0 | 132 | 24.3 | 20.7 |
| 1 | 500 | | | | | | | 505 | | | | 0 | 5 | 1.0 | |
| 3 | 529 | | | | | | | | 619 | | | 0 | 90 | 14.5 | |
| 1 | 845 | | | | | | | | | 792 | | −41 | −12 | −1.5 | |
| 1 | 464 | | | | | | | | | | 563 | −23 | 122 | 21.7 | |
| | Made at 6½ to 7 years old | | | | | | | | | | | | | | |
| 2 | 456 | | 510 | | | | | | | | | 0 | 54 | 10.6 | |
| 12 | 512 | | | 602 | | | | | | | | 0 | 90 | 15.0 | |
| 1 | 499 | | | | 494 | | | | | | | 0 | −5 | −1.0 | |
| 2 | 487 | | | | | 617 | | | | | | 0 | 130 | 12.1 | 14.1 |
| 2 | 470 | | | | | | 510 | | | | | 0 | 40 | 7.8 | |
| 1 | 631 | | | | | | | | 679 | | | 0 | 48 | 7.1 | |
| 1 | 470 | | | | | | | | | 640 | | −23 | 193 | 30.2 | |

TABLE 11.—*Comparison of average initial and retest records of Guernseys having initial records made at ages of 7 and 8 years (single-letter records)*

| Number of cows | Average initial butterfat record made at 7 to 7½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 8 to 8½ | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | 11½ to 12 | 12 to 12½ | 12½ to 13 | 14 to 14½ | | | | |
| | *Pounds* | | | | | | | | | | | *Pounds* | *Pounds* | *Per cent* | *Per cent* |
| 3 | 460 | 538 | | | | | | | | | | 0 | 78 | 14.5 | |
| 3 | 459 | | 643 | | | | | | | | | 0 | 184 | 28.6 | |
| 2 | 496 | | | 594 | | | | | | | | 0 | 98 | 16.5 | |
| 1 | 438 | | | | | | | 490 | | | | −21 | 73 | 14.9 | 21.6 |
| 1 | 485 | | | | | | | | | 702 | | −24 | 241 | 34.3 | |
| 1 | 380 | | | | | | | | | | 490 | −19 | 129 | 26.3 | |
| | Made at 7½ to 8 years old | | | | | | | | | | | | | | |
| 5 | 576 | | 664 | | | | | | | | | 0 | 88 | 13.3 | |
| 1 | 502 | | | 537 | | | | | | | | 0 | 35 | 6.5 | |
| 1 | 371 | | | | 373 | | | | | | | 0 | 2 | 0.5 | 12.6 |
| 1 | 371 | | | | | | 486 | | | | | −18 | 133 | 27.4 | |
| | Made at 8 to 8½ years old | | | | | | | | | | | | | | |
| 1 | 493 | | | 496 | | | | | | | | 0 | 3 | 0.6 | |
| 1 | 501 | | | | 582 | | | | | | | 0 | 81 | 13.9 | 14.5 |
| 1 | 535 | | | | | | | | 717 | | | −26 | 208 | 29.0 | |
| | Made at 8½ to 9 years old | | | | | | | | | | | | | | |
| 2 | 471 | | | 600 | | | | | | | | 0 | 129 | 21.5 | |
| 4 | 501 | | | | 562 | | | | | | | 0 | 61 | 10.9 | |
| 2 | 531 | | | | | 710 | | | | | | 0 | 179 | 25.2 | 17.8 |
| 1 | 572 | | | | | | | 708 | | | | −28 | 164 | 23.2 | |

TABLE 12.—*Comparison of average initial and retest records of Guernseys having initial records made at age of 9 years or older (single-letter records)*

| Number of cows | Average initial butterfat record made at 9 to 9½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 11 to 11½ | 11½ to 12 | 12 to 12½ | 12½ to 13 | 13 to 13½ | 14½ to 15 | | | | |
| | *Pounds* | | | | | | | *Pounds* | *Pounds* | *P. C.* | *Per cent* |
| 1 | 494 | ------ | 488 | ------ | ------ | ------ | ------ | −24 | 18 | 3.7 | 4.8 |
| 1 | 581 | ------ | ------ | 587 | ------ | ------ | ------ | −28 | 34 | 5.8 | |
| | Made at 9½ to 10 years old | | | | | | | | | | |
| 1 | 434 | 481 | ------ | ------ | ------ | ------ | ------ | −21 | 68 | 14.1 | 10.4 |
| 1 | 601 | ------ | ------ | ------ | ------ | 613 | ------ | −29 | 41 | 6.7 | |
| | Made at 10 to 10½ years old | | | | | | | | | | |
| 1 | 392 | ------ | 506 | ------ | ------ | ------ | ------ | −19 | 133 | 26.3 | 26.3 |
| | Made at 11 to 11½ years old | | | | | | | | | | |
| 1 | 433 | ------ | ------ | ------ | 530 | ------ | ------ | 0 | 97 | 18.3 | 18.3 |
| | Made at 12 to 12½ years old | | | | | | | | | | |
| 1 | 410 | ------ | ------ | ------ | ------ | 493 | ------ | 0 | 83 | 16.8 | 16.8 |
| | Made at 13½ to 14 years old | | | | | | | | | | |
| 1 | 609 | ------ | ------ | ------ | ------ | ------ | 713 | 0 | 104 | 14.6 | 14.6 |

TABLE 13.—*Comparison of average initial and retest records of Jerseys having initial records made at age less than 2 years (class A and class AA records)*

| Number of cows | Average initial butterfat record made under 2 years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 2½ to 3 | 3 to 3½ | 3½ to 4 | 4 to 4½ | 4½ to 5 | 5 to 5½ | 5½ to 6 | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 11 to 11½ | | | |
| | *Pounds* | | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *Per cent* | *Per cent* |
| 51 | 340 | 412 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 43 | 29 | 7.0 | |
| 127 | 357 | --- | 452 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 64 | 31 | 6.9 | |
| 40 | 342 | --- | --- | 480 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 80 | 58 | 12.1 | |
| 71 | 343 | --- | --- | --- | 508 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 97 | 68 | 13.4 | |
| 36 | 358 | --- | --- | --- | --- | 525 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 116 | 51 | 9.7 | |
| 28 | 359 | --- | --- | --- | --- | --- | 545 | --- | --- | --- | --- | --- | --- | --- | --- | --- | 119 | 67 | 12.3 | |
| 25 | 365 | --- | --- | --- | --- | --- | --- | 574 | --- | --- | --- | --- | --- | --- | --- | --- | 134 | 75 | 13.1 | |
| 16 | 354 | --- | --- | --- | --- | --- | --- | --- | 551 | --- | --- | --- | --- | --- | --- | --- | 133 | 64 | 11.6 | 10.2 |
| 11 | 387 | --- | --- | --- | --- | --- | --- | --- | --- | 548 | --- | --- | --- | --- | --- | --- | 145 | 16 | 2.9 | |
| 7 | 365 | --- | --- | --- | --- | --- | --- | --- | --- | --- | 607 | --- | --- | --- | --- | --- | 137 | 105 | 17.3 | |
| 8 | 369 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 571 | --- | --- | --- | --- | 139 | 63 | 11.0 | |
| 3 | 336 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 715 | --- | --- | --- | 126 | 253 | 35.4 | |
| 1 | 330 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 785 | --- | --- | 124 | 331 | 42.2 | |
| 2 | 310 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 504 | --- | 117 | 77 | 15.3 | |
| 2 | 328 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 662 | 123 | 211 | 31.9 | |

TABLE 14.—*Comparison of average initial and retest records of Jerseys having initial records made at age of 2 years (class A and class AA)*

| Number of cows | Average initial butterfat record made at 2 to 2½ years old | Average reentry records made at ages shown by same cows as initial records in column at left (pounds of butterfat) | | | | | | | | | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 3 to 3½ | 3½ to 4 | 4 to 4½ | 4½ to 5 | 5 to 5½ | 5½ to 6 | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | 11½ to 12 | 13 to 13½ | 14 to 14½ | 14½ to 15 | 15½ to 16 | | | | |
| | *Pounds* | | | | | | | | | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *P. Ct.* | *Per cent* |
| 240 | 384 | 476 | | | | | | | | | | | | | | | | | | | | | | 37 | 55 | 11.5 | |
| 218 | 395 | | 494 | | | | | | | | | | | | | | | | | | | | | 59 | 40 | 8.1 | |
| 168 | 367 | | | 496 | | | | | | | | | | | | | | | | | | | | 72 | 57 | 11.5 | |
| 152 | 391 | | | | 553 | | | | | | | | | | | | | | | | | | | 92 | 70 | 12.7 | |
| 105 | 388 | | | | | 535 | | | | | | | | | | | | | | | | | | 94 | 53 | 9.9 | |
| 91 | 375 | | | | | | 570 | | | | | | | | | | | | | | | | | 103 | 92 | 16.1 | |
| 64 | 381 | | | | | | | 548 | | | | | | | | | | | | | | | | 108 | 59 | 10.8 | |
| 43 | 382 | | | | | | | | 561 | | | | | | | | | | | | | | | 108 | 71 | 12.7 | |
| 36 | 365 | | | | | | | | | 563 | | | | | | | | | | | | | | 103 | 95 | 16.9 | |
| 35 | 366 | | | | | | | | | | 567 | | | | | | | | | | | | | 104 | 97 | 17.1 | 11.9 |
| 13 | 364 | | | | | | | | | | | 589 | | | | | | | | | | | | 103 | 122 | 20.7 | |
| 12 | 332 | | | | | | | | | | | | 549 | | | | | | | | | | | 94 | 123 | 22.4 | |
| 3 | 322 | | | | | | | | | | | | | 441 | | | | | | | | | | 91 | 28 | 6.4 | |
| 6 | 363 | | | | | | | | | | | | | | 632 | | | | | | | | | 102 | 167 | 26.4 | |
| 2 | 279 | | | | | | | | | | | | | | | 490 | | | | | | | | 79 | 132 | 26.9 | |
| 2 | 393 | | | | | | | | | | | | | | | | | 732 | | | | | | 95 | 244 | 33.3 | |
| 2 | 365 | | | | | | | | | | | | | | | | | | | 528 | | | | 88 | 75 | 14.2 | |
| 1 | 396 | | | | | | | | | | | | | | | | | | | | | 567 | | 57 | 114 | 20.1 | |
| 1 | 396 | | | | | | | | | | | | | | | | | | | | | | 683 | 57 | 230 | 33.7 | |
| | Made at 2½ to 3 years old | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 116 | 395 | | 485 | | | | | | | | | | | | | | | | | | | | | 38 | 52 | 10.7 | |
| 99 | 404 | | | 523 | | | | | | | | | | | | | | | | | | | | 57 | 62 | 11.9 | |
| 53 | 407 | | | | 514 | | | | | | | | | | | | | | | | | | | 72 | 35 | 6.8 | |
| 60 | 399 | | | | | 535 | | | | | | | | | | | | | | | | | | 74 | 62 | 11.6 | |
| 32 | 406 | | | | | | 546 | | | | | | | | | | | | | | | | | 87 | 53 | 9.7 | |
| 43 | 409 | | | | | | | 574 | | | | | | | | | | | | | | | | 91 | 74 | 12.9 | |
| 24 | 387 | | | | | | | | 573 | | | | | | | | | | | | | | | 86 | 100 | 17.5 | |
| 18 | 424 | | | | | | | | | 569 | | | | | | | | | | | | | | 94 | 51 | 9.0 | |
| 15 | 412 | | | | | | | | | | 609 | | | | | | | | | | | | | 92 | 105 | 17.2 | 11.6 |
| 12 | 407 | | | | | | | | | | | 571 | | | | | | | | | | | | 90 | 74 | 13.0 | |
| 8 | 432 | | | | | | | | | | | | 631 | | | | | | | | | | | 96 | 103 | 16.3 | |
| 6 | 399 | | | | | | | | | | | | | 545 | | | | | | | | | | 89 | 57 | 10.5 | |
| 2 | 347 | | | | | | | | | | | | | | 593 | | | | | | | | | 77 | 169 | 28.5 | |
| 3 | 409 | | | | | | | | | | | | | | | 692 | | | | | | | | 91 | 192 | 27.7 | |
| 3 | 407 | | | | | | | | | | | | | | | | 622 | | | | | | | 75 | 140 | 22.5 | |
| 1 | 442 | | | | | | | | | | | | | | | | | 588 | | | | | | 81 | 65 | 11.1 | |
| 1 | 473 | | | | | | | | | | | | | | | | | | 461 | | | | | 87 | −99 | −21.5 | |
| 1 | 442 | | | | | | | | | | | | | | | | | | | | 585 | | | 40 | 103 | 17.6 | |

TABLE 15.—*Comparison of average initial and retest records of Jerseys having initial records made at age of 3 years (class A and class AA records)*

| Number of cows | Average initial butterfat record made at 3 to 3½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 4 to 4½ | 4½ to 5 | 5 to 5½ | 5½ to 6 | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | 12 to 12½ | 12½ to 13 | | | |
| | *Pounds* | | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *P. ct.* |
| 84 | 421 | 498 | | | | | | | | | | | | | | | 37 | 40 | 8.0 | |
| 68 | 434 | | 508 | | | | | | | | | | | | | | 54 | 20 | 3.9 | |
| 56 | 419 | | | 525 | | | | | | | | | | | | | 55 | 51 | 9.7 | |
| 53 | 420 | | | | 518 | | | | | | | | | | | | 68 | 30 | 5.8 | |
| 30 | 421 | | | | | 582 | | | | | | | | | | | 71 | 90 | 15.5 | |
| 29 | 401 | | | | | | 536 | | | | | | | | | | 68 | 67 | 12.5 | |
| 23 | 465 | | | | | | | 603 | | | | | | | | | 78 | 60 | 10.0 | |
| 20 | 420 | | | | | | | | 550 | | | | | | | | 71 | 59 | 10.7 | |
| 15 | 428 | | | | | | | | | 549 | | | | | | | 72 | 49 | 8.9 | 8.7 |
| 7 | 453 | | | | | | | | | | 529 | | | | | | 76 | 0 | 0.0 | |
| 3 | 444 | | | | | | | | | | | 620 | | | | | 75 | 101 | 16.3 | |
| 5 | 421 | | | | | | | | | | | | 542 | | | | 71 | 50 | 9.2 | |
| 8 | 412 | | | | | | | | | | | | | 514 | | | 69 | 33 | 6.4 | |
| 4 | 437 | | | | | | | | | | | | | | 560 | | 57 | 66 | 11.8 | |
| 1 | 437 | | | | | | | | | | | | | | | 555 | 57 | 61 | 11.0 | |
| 2 | 424 | | | | | | | | | | | | | | | | 55 | 230 | 32.4 | |
| | Made at 3½ to 4 years old | | | | | | | | | | | | | | | | | | | |
| 61 | 423 | | 514 | | | | | | | | | | | | | | 31 | 60 | 11.7 | |
| 57 | 446 | | | 532 | | | | | | | | | | | | | 36 | 50 | 9.4 | |
| 33 | 431 | | | | 519 | | | | | | | | | | | | 47 | 41 | 7.9 | |
| 41 | 416 | | | | | 495 | | | | | | | | | | | 48 | 31 | 6.3 | |
| 16 | 446 | | | | | | 550 | | | | | | | | | | 51 | 53 | 9.6 | |
| 14 | 432 | | | | | | | 581 | | | | | | | | | 50 | 99 | 17.0 | |
| 11 | 374 | | | | | | | | 508 | | | | | | | | 43 | 91 | 17.9 | |
| 7 | 407 | | | | | | | | | 560 | | | | | | | 47 | 106 | 18.9 | 11.5 |
| 11 | 409 | | | | | | | | | | 618 | | | | | | 47 | 162 | 26.2 | |
| 5 | 422 | | | | | | | | | | | 547 | | | | | 49 | 76 | 13.9 | |
| 3 | 430 | | | | | | | | | | | | 678 | | | | 50 | 198 | 29.2 | |
| 2 | 460 | | | | | | | | | | | | | 641 | | | 53 | 128 | 20.0 | |
| 3 | 440 | | | | | | | | | | | | | | 548 | | 35 | 73 | 13.3 | |
| 1 | 512 | | | | | | | | | | | | | | | 741 | 41 | 188 | 25.4 | |
| 2 | 515 | | | | | | | | | | | | | | | | 41 | 109 | 16.4 | |

TABLE 16.—*Comparison of average initial and retest records of Jerseys having initial records made at age of 4 years (class A and class AA records)*

| Number of cows | Average initial butterfat record made at 4 to 4½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 5 to 5½ | 5½ to 6 | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | 11½ to 12 | 12½ to 13 | | | |
| | *Pounds* | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *Per cent* |
| 66 | 436 | 488 | | | | | | | | | | | | | | 17 | 35 | 7.2 | |
| 53 | 453 | | 540 | | | | | | | | | | | | | 30 | 57 | 10.6 | |
| 19 | 436 | | | 502 | | | | | | | | | | | | 32 | 34 | 6.8 | |
| 21 | 444 | | | | 526 | | | | | | | | | | | 32 | 50 | 9.5 | |
| 23 | 470 | | | | | 574 | | | | | | | | | | 34 | 70 | 12.2 | |
| 12 | 424 | | | | | | 569 | | | | | | | | | 31 | 114 | 20.0 | |
| 11 | 477 | | | | | | | 600 | | | | | | | | 35 | 88 | 14.7 | |
| 7 | 512 | | | | | | | | 642 | | | | | | | 37 | 93 | 14.5 | 10.3 |
| 3 | 454 | | | | | | | | | 555 | | | | | | 33 | 68 | 12.3 | |
| 2 | 458 | | | | | | | | | | 540 | | | | | 33 | 49 | 9.1 | |
| 2 | 419 | | | | | | | | | | | 512 | | | | 30 | 63 | 12.3 | |
| 2 | 368 | | | | | | | | | | | | 408 | | | 14 | 26 | 6.4 | |
| 1 | 389 | | | | | | | | | | | | | 589 | | 15 | 185 | 31.4 | |
| 2 | 458 | | | | | | | | | | | | | | 548 | 18 | 72 | 13.1 | |
| | Made at 4½ to 5 years old | | | | | | | | | | | | | | | | | | |
| 39 | 449 | | 513 | | | | | | | | | | | | | 15 | 49 | 9.6 | |
| 29 | 479 | | | 567 | | | | | | | | | | | | 19 | 69 | 12.2 | |
| 22 | 468 | | | | 528 | | | | | | | | | | | 19 | 41 | 7.8 | |
| 19 | 435 | | | | | 515 | | | | | | | | | | 19 | 63 | 12.2 | |
| 4 | 519 | | | | | | 531 | | | | | | | | | 20 | −8 | −1.5 | |
| 12 | 466 | | | | | | | 558 | | | | | | | | 18 | 74 | 13.3 | |
| 4 | 464 | | | | | | | | 706 | | | | | | | 18 | 224 | 31.7 | 11.5 |
| 9 | 416 | | | | | | | | | 557 | | | | | | 16 | 125 | 22.4 | |
| 5 | 485 | | | | | | | | | | 630 | | | | | 19 | 126 | 20.0 | |
| 1 | 535 | | | | | | | | | | | 406 | | | | 21 | −150 | −36.9 | |
| 1 | 537 | | | | | | | | | | | | 488 | | | 4 | −53 | −10.9 | |
| 2 | 443 | | | | | | | | | | | | | 539 | | 3 | 93 | 17.3 | |
| 1 | 437 | | | | | | | | | | | | | | 457 | 3 | 17 | 3.7 | |

TABLE 17.—*Comparison of average initial and retest records of Jerseys having initial records made at age of 5 years (class A and class AA records)*

| Number of cows | Average initial butterfat record made at 5 to 5½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 6 to 6½ | 6½ to 7 | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | 11½ to 12 | 12½ to 13 | 14 to 14½ | | | |
| | *Pounds* | | | | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *Per cent* |
| 26 | 456 | 508 | | | | | | | | | | | | | | 15 | 37 | 7.3 | |
| 35 | 477 | | 556 | | | | | | | | | | | | | 16 | 63 | 11.3 | |
| 17 | 452 | | | 537 | | | | | | | | | | | | 15 | 70 | 13.0 | |
| 12 | 497 | | | | 553 | | | | | | | | | | | 17 | 39 | 7.1 | |
| 8 | 434 | | | | | 575 | | | | | | | | | | 14 | 127 | 22.1 | |
| 8 | 451 | | | | | | 543 | | | | | | | | | 15 | 77 | 14.2 | 11.7 |
| 9 | 532 | | | | | | | 667 | | | | | | | | 17 | 118 | 17.7 | |
| 3 | 444 | | | | | | | | 554 | | | | | | | 15 | 95 | 17.1 | |
| 2 | 422 | | | | | | | | | 410 | | | | | | 14 | −26 | −6.3 | |
| 1 | 498 | | | | | | | | | | 588 | | | | | 0 | 90 | 15.3 | |
| 1 | 384 | | | | | | | | | | | | | 529 | | 0 | 145 | 27.4 | |
| | Made at 5½ to 6 years old | | | | | | | | | | | | | | | | | | |
| 36 | 458 | | 520 | | | | | | | | | | | | | 3 | 59 | 11.3 | |
| 24 | 510 | | | 573 | | | | | | | | | | | | 3 | 60 | 10.5 | |
| 20 | 467 | | | | 552 | | | | | | | | | | | 3 | 82 | 14.9 | |
| 19 | 495 | | | | | 567 | | | | | | | | | | 3 | 69 | 12.2 | |
| 4 | 452 | | | | | | 485 | | | | | | | | | 3 | 30 | 6.2 | |
| 4 | 536 | | | | | | | 570 | | | | | | | | 3 | 31 | 5.4 | |
| 5 | 508 | | | | | | | | 499 | | | | | | | 3 | −12 | −2.4 | 11.1 |
| 1 | 536 | | | | | | | | | 494 | | | | | | 3 | −45 | −9.1 | |
| 3 | 503 | | | | | | | | | | 542 | | | | | −13 | 52 | 9.6 | |
| 2 | 531 | | | | | | | | | | | 661 | | | | −14 | 144 | 21.8 | |
| 2 | 429 | | | | | | | | | | | | 608 | | | −11 | 190 | 31.3 | |
| 3 | 495 | | | | | | | | | | | | | 471 | | −13 | −11 | −2.3 | |
| 1 | 361 | | | | | | | | | | | | | | 506 | −37 | 182 | 36.0 | |

TABLE 18.—*Comparison of average initial and retest records of Jerseys having initial records made at age of 6 years (class A and class AA records)*

| Number of cows | Average initial butterfat record made at 6 to 6½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 7 to 7½ | 7½ to 8 | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 11 to 11½ | 11½ to 12 | 12 to 12½ | 15 to 15½ | | | | |
| | *Pounds* | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *Per cent* |
| 34 | 494 | 549 | | | | | | | | | | | 0 | 55 | 10.0 | |
| 23 | 523 | | 548 | | | | | | | | | | 0 | 25 | 4.6 | |
| 10 | 520 | | | 565 | | | | | | | | | 0 | 45 | 8.0 | |
| 10 | 453 | | | | 517 | | | | | | | | 0 | 64 | 12.4 | |
| 4 | 472 | | | | | 557 | | | | | | | 0 | 85 | 15.3 | 9.8 |
| 3 | 464 | | | | | | 562 | | | | | | 0 | 98 | 17.4 | |
| 1 | 415 | | | | | | | 724 | | | | | 0 | 309 | 42.7 | |
| 1 | 408 | | | | | | | | 376 | | | | −13 | −19 | −5.1 | |
| 4 | 509 | | | | | | | | | 695 | | | −16 | 202 | 29.1 | |
| 1 | 604 | | | | | | | | | | 452 | | −19 | −133 | −29.4 | |
| | Made at 6½ to 7 years old | | | | | | | | | | | | | | | |
| 25 | 452 | | 519 | | | | | | | | | | 0 | 67 | 12.9 | |
| 14 | 563 | | | 591 | | | | | | | | | 0 | 28 | 4.7 | |
| 2 | 563 | | | | 530 | | | | | | | | 0 | −33 | −6.2 | |
| 8 | 518 | | | | | 560 | | | | | | | 0 | 42 | 7.5 | |
| 2 | 423 | | | | | | 416 | | | | | | 0 | −7 | −1.7 | 8.0 |
| 1 | 473 | | | | | | | 508 | | | | | 0 | 35 | 6.9 | |
| 1 | 594 | | | | | | | | 852 | | | | −19 | 277 | 32.5 | |
| 1 | 425 | | | | | | | | | 381 | | | −14 | −30 | −7.9 | |
| 1 | 529 | | | | | | | | | | | 502 | −57 | 30 | 6.0 | |

TABLE 19.—*Comparison of average initial and retest records of Jerseys having initial records made at ages of 7 and 8 years (class A and class AA records)*

| Number of cows | Average initial butterfat record made at 7 to 7½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 8 to 8½ | 8½ to 9 | 9 to 9½ | 9½ to 10 | 10 to 10½ | 10½ to 11 | 11 to 11½ | 11½ to 12 | 12 to 12½ | 12½ to 13 | 13 to 13½ | 13½ to 14 | 14½ to 15 | | | |
| | *Pounds* | | | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *Per cent* |
| 19 | 488 | 517 | | | | | | | | | | | | | 0 | 29 | 5.6 | |
| 14 | 519 | | 559 | | | | | | | | | | | | 0 | 40 | 7.1 | |
| 3 | 559 | | | 488 | | | | | | | | | | | 0 | −71 | −14.5 | |
| 10 | 471 | | | | 492 | | | | | | | | | | 0 | 21 | 4.3 | 6.3 |
| 2 | 434 | | | | | 462 | | | | | | | | | 0 | 28 | 6.1 | |
| 2 | 433 | | | | | | 573 | | | | | | | | −14 | 154 | 26.9 | |
| 1 | 394 | | | | | | | | | | | | | 683 | −42 | 331 | 48.5 | |
| | Made at 7½ to 8 years old | | | | | | | | | | | | | | | | | |
| 6 | 468 | | 580 | | | | | | | | | | | | 0 | 112 | 19.3 | |
| 10 | 501 | | | 540 | | | | | | | | | | | 0 | 39 | 7.2 | |
| 1 | 433 | | | | 510 | | | | | | | | | | 0 | 77 | 15.1 | |
| 1 | 500 | | | | | 457 | | | | | | | | | 0 | −43 | −9.4 | |
| 1 | 493 | | | | | | 591 | | | | | | | | −16 | 114 | 19.3 | 13.8 |
| 1 | 487 | | | | | | | | 829 | | | | | | −15 | 357 | 43.1 | |
| 2 | 485 | | | | | | | | | | 620 | | | | −15 | 150 | 24.2 | |
| 1 | 502 | | | | | | | | | | | 555 | | | −16 | 69 | 12.4 | |
| | Made at 8 to 8½ years old | | | | | | | | | | | | | | | | | |
| 10 | 477 | | | 534 | | | | | | | | | | | 0 | 57 | 10.7 | |
| 11 | 523 | | | | 556 | | | | | | | | | | 0 | 33 | 5.9 | |
| 5 | 512 | | | | | 565 | | | | | | | | | 0 | 53 | 9.4 | |
| 2 | 463 | | | | | | 501 | | | | | | | | −15 | 53 | 10.6 | 8.1 |
| 3 | 518 | | | | | | | 535 | | | | | | | −17 | 34 | 6.4 | |
| 1 | 432 | | | | | | | | 663 | | | | | | −14 | 245 | 37.0 | |
| 1 | 531 | | | | | | | | | | | | 397 | | −17 | −117 | −29.5 | |
| | Made at 8½ to 9 years old | | | | | | | | | | | | | | | | | |
| 6 | 429 | | | | 504 | | | | | | | | | | 0 | 75 | 14.9 | |
| 1 | 491 | | | | | 485 | | | | | | | | | 0 | −6 | −1.2 | |
| 2 | 383 | | | | | | 596 | | | | | | | | −12 | 225 | 37.8 | |
| 2 | 531 | | | | | | | | 547 | | | | | | −17 | 33 | 6.0 | 14.8 |
| 1 | 398 | | | | | | | | | 383 | | | | | −13 | −2 | −0.5 | |
| 1 | 389 | | | | | | | | | | 456 | | | | −12 | 79 | 17.3 | |

TABLE 20.—*Comparison of average initial and retest records of Jerseys having initial records made at ages of 9 and 10 years (class A and class AA records)*

| Number of cows | Average initial butterfat record made at 9 to 9½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 10 to 10½ | 10½ to 11 | 11 to 11½ | 11½ to 12 | 12 to 12½ | 12½ to 13 | 13½ to 14 | 14½ to 15 | | | | |
| | *Pounds* | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *Per cent* |
| 7 | 450 | 516 | | | | | | | | 0 | 66 | 12.8 | 13.8 |
| 3 | 480 | | 657 | | | | | | | −15 | 192 | 29.2 | |
| 1 | 407 | | | 425 | | | | | | −13 | 31 | 7.3 | |
| 1 | 471 | | | | 382 | | | | | −15 | −74 | −19.4 | |
| | Made at 9½ to 10 years old | | | | | | | | | | | | |
| 2 | 652 | | 741 | | | | | | | −21 | 110 | 14.8 | 5.4 |
| 1 | 495 | | | 539 | | | | | | −16 | 60 | 11.1 | |
| 1 | 385 | | | | 395 | | | | | −12 | 22 | 5.6 | |
| 1 | 495 | | | | | 404 | | | | −16 | −75 | −18.6 | |
| 1 | 495 | | | | | | | 503 | | −16 | 24 | 4.8 | |
| | Made at 10 to 10½ years old | | | | | | | | | | | | |
| 7 | 484 | | | 483 | | | | | | −16 | 15 | 3.1 | 7.4 |
| 1 | 426 | | | | 425 | | | | | −13 | 12 | 2.8 | |
| 2 | 392 | | | | | 500 | | | | −12 | 120 | 24.0 | |
| 2 | 430 | | | | | | 452 | | | −14 | 36 | 8.0 | |
| | Made at 10½ to 11 years old | | | | | | | | | | | | |
| 3 | 518 | | | | 461 | | | | | 0 | −57 | −12.4 | −5.8 |
| 1 | 501 | | | | | 478 | | | | 0 | −23 | −4.8 | |
| 1 | 573 | | | | | | | | 606 | −45 | 78 | 12.9 | |

TABLE 21.—*Comparison of average initial and retest records of Jerseys having initial records made at age of 11 years or over (class A and class AA records)*

| Number of cows | Average initial butterfat record made at 11 to 11½ years old | Average reentry records made at ages shown in these columns by same cows as initial records shown in column at left (pounds of butterfat) | | | | | | | | | | Increase due to age | Portion of reentry record due to development | | Average due to development for this group |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 12 to 12½ | 12½ to 13 | 13 to 13½ | 14 to 14½ | 14½ to 15 | 15 to 15½ | 16 to 16½ | 16½ to 17 | 17½ to 18 | 18 to 18½ | 19 to 19½ | | | |
| | *Pounds* | | | | | | | | | | | | *Pounds* | *Pounds* | *P. ct.* | *Per cent* |
| 2 | 395 | 430 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 0 | 35 | 8.1 | 8.1 |
| | Made at 11½ to 12 years old | | | | | | | | | | | | | | | |
| 3 | 417 | --- | 421 | --- | --- | --- | --- | --- | --- | --- | --- | --- | 0 | 4 | 1.0 | 1.0 |
| | Made at 12 to 12½ years old | | | | | | | | | | | | | | | |
| 2 | 475 | --- | --- | 512 | --- | --- | --- | --- | --- | --- | --- | --- | 0 | 37 | 7.2 | 5.5 |
| 1 | 441 | --- | --- | --- | --- | --- | --- | --- | --- | --- | 413 | --- | −37 | 9 | 2.2 | --- |
| | Made at 13 to 13½ years old | | | | | | | | | | | | | | | |
| 1 | 482 | --- | --- | --- | 431 | --- | --- | --- | --- | --- | --- | --- | −38 | −13 | −3.0 | --- |
| 1 | 364 | --- | --- | --- | --- | 509 | --- | --- | --- | --- | --- | --- | −28 | 173 | 34.0 | 13.1 |
| 1 | 476 | --- | --- | --- | --- | --- | 479 | --- | --- | --- | --- | --- | −37 | 40 | 8.4 | --- |
| | Made at 13½ to 14 years old | | | | | | | | | | | | | | | |
| 1 | 483 | --- | --- | --- | --- | --- | 579 | --- | --- | --- | --- | --- | −38 | 134 | 23.1 | 19.0 |
| 1 | 483 | --- | --- | --- | --- | --- | --- | 523 | --- | --- | --- | --- | −38 | 78 | 14.9 | --- |
| | Made at 14 to 14½ years old | | | | | | | | | | | | | | | |
| 1 | 487 | --- | --- | --- | --- | --- | 523 | --- | --- | --- | --- | --- | 0 | 36 | 6.9 | --- |
| 1 | 487 | --- | --- | --- | --- | --- | --- | --- | 530 | --- | --- | --- | 0 | 43 | 8.1 | 10.8 |
| 1 | 487 | --- | --- | --- | --- | --- | --- | --- | --- | 590 | --- | --- | 0 | 103 | 17.5 | --- |
| | Made at 18 to 18½ years old | | | | | | | | | | | | | | | |
| 1 | 401 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 401 | 0 | 0 | 0.0 | 0.0 |

It is a natural assumption that there is a rather high correlation between the initial and reentry records of the same cow or group of cows. Tables 23 and 22 give the distribution of the 240 Jerseys having initial records made at 2 to 2½ years of age and reentry records made when 3 to 3½ years old, and of the 106 Guernsey cows having initial records made when 2 to 2½ years old and reentry records made at 3½ to 4 years of age. The coefficient of correlation of these Jersey records is $0.6318 \pm 0.0262$, and of the Guernseys, $0.7247 \pm 0.0311$. These two groups were selected because they contain the largest number of individuals of all of the groups of their respective breeds.

TABLE 22.—*Distribution of initial and reentry records of Guernsey cows, initial records made at 2 to 2½ years of age and reentry records at 3½ to 4 years of age*

| Initial butterfat records (pounds) | Reentry records (pounds of butterfat) | | | | | | | | | | | | | | | | | | | | | | Total |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 350 to 374 | 375 to 399 | 400 to 424 | 425 to 449 | 450 to 474 | 475 to 499 | 500 to 524 | 525 to 549 | 550 to 574 | 575 to 599 | 600 to 624 | 625 to 649 | 650 to 674 | 675 to 699 | 700 to 724 | 725 to 749 | 750 to 774 | 775 to 799 | 800 to 824 | 825 to 849 | 850 to 874 | 875 to 899 | 900 to 924 | |
| 250 to 274 | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 |
| 275 to 299 | 1 | 2 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 3 |
| 300 to 324 | --- | --- | 1 | --- | --- | --- | --- | --- | --- | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 2 |
| 325 to 349 | 1 | 1 | 2 | 2 | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 7 |
| 350 to 374 | --- | 2 | 1 | 2 | --- | 1 | 1 | --- | --- | 1 | 1 | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 10 |
| 375 to 399 | --- | 1 | 4 | 1 | 2 | --- | 1 | --- | 1 | 1 | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 12 |
| 400 to 424 | --- | --- | --- | 2 | 2 | 3 | 2 | 2 | --- | 1 | --- | 1 | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 14 |
| 425 to 449 | --- | --- | 1 | 1 | 1 | 1 | 2 | --- | 1 | 1 | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 9 |
| 450 to 474 | --- | --- | --- | --- | --- | --- | 1 | 1 | 3 | 3 | --- | --- | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | 9 |
| 475 to 499 | --- | --- | --- | --- | 2 | 1 | 2 | --- | --- | 2 | 2 | 1 | --- | 1 | --- | --- | --- | 1 | --- | --- | --- | --- | --- | 12 |
| 500 to 524 | --- | --- | --- | --- | --- | 1 | --- | 1 | 2 | --- | 1 | 3 | 3 | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | 12 |
| 525 to 549 | --- | --- | --- | --- | --- | --- | 1 | --- | --- | --- | --- | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 2 |
| 550 to 574 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | 1 | --- | 1 | --- | --- | --- | 2 | --- | --- | --- | --- | 5 |
| 600 to 624 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | 1 | --- | --- | --- | 1 | --- | --- | --- | --- | 3 |
| 625 to 649 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | 1 | --- | --- | --- | --- | --- | 2 |
| 650 to 674 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 |
| 675 to 699 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 700 to 724 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | 1 |
| 725 to 749 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | --- | --- | --- | --- | --- | --- | 1 |
| Total | 2 | 7 | 9 | 8 | 7 | 8 | 9 | 5 | 7 | 9 | 5 | 8 | 7 | 4 | 3 | --- | 2 | 2 | 3 | --- | --- | --- | 1 | 106 |

TABLE 23.—*Distribution of initial and reentry records of Jersey cows, initial records made at 2 to 2½ years of age and reentry records at 3 to 3½ years of age*

| Initial butterfat records (pounds) | Reentry records (pounds of butterfat) | | | | | | | | | | | | | | | | | | | | | | | Total |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | 300 to 324 | 325 to 349 | 350 to 374 | 375 to 399 | 400 to 424 | 425 to 449 | 450 to 474 | 475 to 499 | 500 to 524 | 525 to 549 | 550 to 574 | 575 to 599 | 600 to 624 | 625 to 649 | 650 to 674 | 675 to 699 | 700 to 724 | 725 to 749 | 750 to 774 | 775 to 799 | 800 to 824 | 825 to 849 | 850 to 874 | 875 to 899 | |
| 250 to 274 | 4 | 5 | 1 | --- | 3 | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 14 |
| 275 to 299 | 3 | 1 | 2 | --- | --- | --- | 5 | 1 | --- | --- | --- | 1 | 1 | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 15 |
| 300 to 324 | 1 | 8 | 3 | 8 | 5 | 4 | --- | 2 | --- | 1 | 1 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 33 |
| 325 to 349 | --- | 3 | 5 | 6 | 4 | 3 | 3 | 2 | 1 | 2 | 2 | 1 | 1 | --- | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | 34 |
| 350 to 374 | 3 | 1 | 2 | 4 | 1 | 4 | 5 | 1 | 1 | 2 | 2 | 2 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 28 |
| 375 to 399 | 1 | --- | 2 | 2 | 4 | 2 | 2 | 7 | 3 | 3 | 3 | 1 | 2 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 32 |
| 400 to 424 | --- | --- | --- | --- | 2 | 3 | 2 | 4 | 4 | 1 | 2 | 2 | --- | --- | 1 | --- | 1 | --- | --- | --- | --- | --- | --- | --- | 22 |
| 425 to 449 | 1 | 1 | --- | --- | --- | --- | 3 | 2 | 2 | --- | --- | 2 | --- | 3 | 1 | 1 | --- | 2 | --- | --- | --- | --- | --- | --- | 18 |
| 450 to 474 | --- | 1 | --- | --- | --- | 1 | 1 | 1 | --- | 2 | 1 | 1 | --- | --- | --- | 2 | --- | --- | 1 | --- | --- | --- | --- | --- | 11 |
| 475 to 499 | --- | 1 | 1 | --- | 2 | 1 | --- | --- | --- | --- | 2 | 1 | --- | --- | --- | --- | --- | --- | 1 | --- | --- | --- | --- | --- | 9 |
| 500 to 524 | --- | --- | --- | --- | --- | --- | --- | --- | 1 | 1 | 1 | 1 | --- | --- | --- | 1 | --- | --- | --- | --- | --- | --- | --- | --- | 5 |
| 525 to 549 | --- | --- | --- | --- | --- | --- | --- | 1 | 2 | --- | --- | 1 | 1 | 1 | --- | --- | 2 | --- | --- | --- | --- | --- | --- | --- | 8 |
| 550 to 574 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | 1 | --- | --- | --- | --- | 1 | --- | --- | 1 | --- | --- | --- | --- | 4 |
| 575 to 599 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | --- | --- | 1 | 1 | --- | --- | --- | --- | --- | --- | --- | 3 |
| 600 to 624 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | --- | --- | --- | --- | 1 | --- | --- | --- | --- | --- | --- | --- | 2 |
| 625 to 649 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | 1 |
| 650 to 674 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 |
| 675 to 699 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 700 to 724 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 725 to 749 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 750 to 774 | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | 1 | --- | --- |
| Total | 13 | 21 | 16 | 20 | 21 | 19 | 21 | 21 | 13 | 12 | 15 | 15 | 7 | 5 | 2 | 6 | 6 | 2 | 2 | 1 | --- | 1 | --- | 1 | 240 |

The following reasons are usually given for retesting cows:

(1) Those that have shown unusual ability during the first test are retested with the hope of making a still more creditable record.

(2) The cow of good breeding whose initial record did not come up to expectations is retested with the hope that on the next test she

will show improvement, not only for her own sake, but also to add prestige to the family of which she is a member.

(3) The cow that failed to come up to expectations on her first test because of injury or some other known reason is retested in order to obtain a normal and better record.

Tables 24 and 25 were prepared in order to determine whether the cows retested were better or poorer producers on the average than all tested cows.

TABLE 24.—*Average initial butterfat records of all advanced-registry Guernsey cows and those of cows retested (classified according to age at time of initial test)*

| Age | Initial records of all cows | | Initial records of cows retested | | Age | Initial records of all cows | | Initial records of cows retested | |
|---|---|---|---|---|---|---|---|---|---|
| | Number | Butterfat | Number | Butterfat | | Number | Butterfat | Number | Butterfat |
| | | *Pounds* | | *Pounds* | | | *Pounds* | | *Pounds* |
| Under 2 years | 275 | 368 | 55 | 381 | 7 to 7½ years | 179 | 495 | 11 | 465 |
| 2 to 2½ years | 2,367 | 404 | 449 | 427 | 7½ to 8 years | 148 | 499 | 8 | 516 |
| 2½ to 3 years | 1,211 | 424 | 215 | 441 | 8 to 8½ years | 135 | 496 | 3 | 510 |
| 3 to 3½ years | 897 | 433 | 146 | 443 | 8½ to 9 years | 88 | 495 | 9 | 509 |
| 3½ to 4 years | 695 | 453 | 99 | 463 | 9 to 9½ years | 102 | 502 | 2 | 538 |
| 4 to 4½ years | 625 | 470 | 76 | 473 | 9½ to 10 years | 80 | 496 | 2 | 518 |
| 4½ to 5 years | 432 | 473 | 42 | 494 | 10 to 10½ years | 45 | 493 | 1 | 392 |
| 5 to 5½ years | 373 | 495 | 22 | 506 | 11 to 11½ years | 28 | 459 | 1 | 433 |
| 5½ to 6 years | 313 | 489 | 23 | 504 | 12 to 12½ years | 13 | 504 | 1 | 410 |
| 6 to 6½ years | 305 | 482 | 28 | 499 | 13½ to 14 years | 8 | 494 | 1 | 609 |
| 6½ to 7 years | 214 | 487 | 21 | 503 | | | | | |

TABLE 25.—*Average initial butterfat records of all register-of-merit Jersey cows and those of cows retested (classified according to age at time of initial test)*

| Age | Initial records of all cows | | Initial records of cows retested | | Age | Initial records of all cows | | Initial records of cows retested | |
|---|---|---|---|---|---|---|---|---|---|
| | Number | Butterfat | Number | Butterfat | | Number | Butterfat | Number | Butterfat |
| | | *Pounds* | | *Pounds* | | | *Pounds* | | *Pounds* |
| Under 2 years | 1,237 | 343 | 428 | 352 | 8 to 8½ years | 267 | 480 | 33 | 501 |
| 2 to 2½ years | 3,633 | 368 | 1,194 | 382 | 8½ to 9 years | 232 | 462 | 13 | 437 |
| 2½ to 3 years | 1,613 | 386 | 497 | 403 | 9 to 9½ years | 163 | 468 | 12 | 456 |
| 3 to 3½ years | 1,398 | 404 | 408 | 425 | 9½ to 10 years | 128 | 475 | 6 | 529 |
| 3½ to 4 years | 1,011 | 423 | 267 | 428 | 10 to 10½ years | 114 | 476 | 12 | 455 |
| 4 to 4½ years | 923 | 440 | 224 | 448 | 10½ to 11 years | 68 | 459 | 5 | 526 |
| 4½ to 5 years | 705 | 454 | 148 | 460 | 11 to 11½ years | 53 | 454 | 2 | 395 |
| 5 to 5½ years | 673 | 457 | 122 | 468 | 11½ to 12 years | 39 | 478 | 3 | 417 |
| 5½ to 6 years | 587 | 469 | 124 | 478 | 12 to 12½ years | 40 | 445 | 3 | 463 |
| 6 to 6½ years | 509 | 475 | 91 | 498 | 13 to 13½ years | 18 | 445 | 3 | 441 |
| 6½ to 7 years | 404 | 471 | 55 | 497 | 13½ to 14 years | 11 | 475 | 2 | 483 |
| 7 to 7½ years | 395 | 468 | 51 | 491 | 14 to 14½ years | 6 | 439 | 3 | 487 |
| 7½ to 8 years | 310 | 470 | 23 | 487 | | | | | |

Except in some of the older classes having only a small number of individuals, the animals retested had initial records above the average for all cows in the group of that age. This fact has no bearing on the information given in Tables 5 to 21, as allowance is made for this difference between the initial records of the animals under consideration and the average of all initial records, and the difference is carried into the factor representing the increase due to age. The factor of development is not affected.

## EFFECT OF PREGNANCY

That pregnancy has a pronounced influence on the quantity of butterfat produced during a year when the calf is carried for a period of 265 days or longer is evidenced by the average of 408 pounds of butterfat for 711 class GG records as compared with an average of 418 pounds of butterfat for 4,794 class G records of the Guernsey breed. All these, of course, are initial records. The other classes of Guernseys were not studied.

The register-of-merit rules specified that a Jersey cow carrying a calf for 155 days during the period of her test was entitled to entry in class AA. One feature about the class AA Jersey records which tends to nullify the value of any conclusion drawn therefrom, is the fact that in the past many breeders failed to apply for transfer of animals from class A to class AA after they had fulfilled the calving requirement, and consequently there is a certain proportion of the animals which rightly belong to class AA now listed in the other class.

For this reason, and in order to enhance the value of the class averages by using greater numbers, it was thought best to deal with all Jersey records as a single class. However, classes A and AA were compiled separately, and it was found that even with the moderate demands of the former Jersey rules some influence of gestation is apparent in the records. It is well to remember that the group of class AA cows includes some animals that carried a calf full time during the test, and that they grade down to those which just met the specified 155-day period. It might be assumed that the average time that these class AA cows were pregnant during their test periods would be a somewhere near 200 days.

For purposes of comparison, Table 26 lists the Jersey class A and class AA records in parallel columns.

TABLE 26.—*Initial butterfat records for class A and class AA Jersey cows*

| Age | Class A | | Class AA | | Difference in butterfet | Age | Class A | | Class AA | | Difference in butterfet |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | Records | Production of butterfat | Records | Production of butterfat | | | Records | Production of butterfat | Records | Production of butterfat | |
| | *No.* | *Lbs.* | *No.* | *Lbs.* | *Lbs.* | | *No.* | *Lbs.* | *No.* | *Lbs.* | *Lbs.* |
| Under 2 years | 461 | 344 | 776 | 342 | 2 | 8 to 8½ years | 105 | 497 | 162 | 469 | 28 |
| 2 to 2½ years | 1,567 | 368 | 2,066 | 368 | 0 | 8½ to 9 years | 104 | 465 | 128 | 460 | 5 |
| 2½ to 3 years | 669 | 386 | 944 | 387 | 1 | 9 to 9½ years | 77 | 485 | 86 | 454 | 31 |
| 3 to 3½ years | 586 | 409 | 812 | 400 | 9 | 9½ to 10 years | 67 | 483 | 61 | 467 | 16 |
| 3½ to 4 years | 385 | 427 | 626 | 420 | 7 | 10 to 10½ years | 50 | 487 | 64 | 468 | 19 |
| 4 to 4½ years | 350 | 446 | 574 | 437 | 9 | 10½ to 11 years | 28 | 459 | 40 | 459 | 0 |
| 4½ to 5 years | 283 | 457 | 422 | 451 | 6 | 11 to 11½ years | 26 | 468 | 27 | 440 | 28 |
| 5 to 5½ years | 265 | 465 | 408 | 453 | 12 | 11½ to 12 years | 17 | 499 | 22 | 461 | 38 |
| 5½ to 6 years | 219 | 476 | 368 | 465 | 11 | 12 to 12½ years | 16 | 452 | 24 | 440 | 12 |
| 6 to 6½ years | 209 | 482 | 300 | 470 | 12 | 12½ to 13 years | 13 | 438 | 3 | 427 | 11 |
| 6½ to 7 years | 165 | 478 | 239 | 462 | 16 | 13 to 13½ years | 8 | 472 | 10 | 424 | 48 |
| 7 to 7½ years | 168 | 482 | 227 | 458 | 24 | 13½ to 14 years | 6 | 495 | 5 | 451 | 44 |
| 7½ to 8 years | 134 | 480 | 176 | 463 | 17 | | | | | | |

Where the groups contain a considerable number of animals the difference in pounds of butterfat is not large enough to be of much significance, though evidence of the effect of gestation is found in the fact that every group, with the exception of those 2 years old,

shows a difference in favor of the class A animals, or those which did not meet the calving requirement of the rules. As before mentioned, there may be some cows in the single-letter class which did carry calves 155 days or more but, through neglect of the owner, were not changed to class AA.

Another significant point brought out by this table is that the tax of pregnancy seems to grow more severe with advancing age. The difference between the two classes for all ages up to 3 years is insignificant, but from that age on there is a marked divergence between the average records for the two classes; and it would appear that, while the young animal carries on the function of reproduction without its affecting materially her productive capacity, the older cows do not stand up so well under this double strain.

## RECENT IMPROVEMENT IN PRODUCTION

An interesting fact brought out incidentally by this study is the noticeable improvement in production records since the beginning of 1920, especially in the last thousand records. This is illustrated in Figure 1. All the animals shown are classed according to age; and in each age class an average record is computed for each successive group of 1,000 records in the Register of Merit. These averages are plotted in solid lines, and the dotted lines represent the average record for each age group over the entire period of time. At the top of the graph is shown the year in which each block of 1,000 records was completed.

The chart shows that there has been a general improvement in all classes, most of which appears in the last 6,000 records. The rise is gradual and would seem to indicate that with the increase in experience gained by handling test cows the breeders have developed better methods of feeding, conditioning, and caring for animals on test which have brought about better average records. The effect of years of selective breeding for high production is also a force working to raise these averages.

That some such factors have raised the general average of all records also helps to account for the fact that the reentry records made after a lapse of six years or more usually show a larger increase due to development than those made after a shorter interval. We have here a combination of development of the cows and methods of caring for them.

## SUMMARY AND CONCLUSIONS

Initial advanced-registry Guernsey records and register-of-merit Jersey records furnished material for Tables 1 and 2, which show the average production for animals grouped according to age.

A study of these tables shows the actual relationship of average records made at different ages.

The results will be used as a basis for adjusting records of various ages in future studies of prepotency of sires made by this bureau.

The age of maximum production for the Jersey cows included in this study begins when the cow is 6 years old and continues until she reaches the age of 10½ years.

The data on the Guernsey cows studied indicate that they reach their maximum producing ability at 5 years of age, and the period of maximum production continues until they are 11 years old.

The results of this study do not furnish a basis for interbreed comparisons, since it embodies Jersey records with and without calving requirements; whereas the Guernsey records were all of the




FIG. 1.—Improvement in butterfat production of Jersey cows, as shown by the average production of all records for each class compared with the average production of each thousand records, 1911 to 1921. Dotted line: Average production for entire period. Solid line: Average production by groups of 1,000 records

single-letter class. Furthermore, no information is given to indicate what proportion of the cows were milked more than twice daily.

The increase in production from an initial record to a reentry record is owing partly to increased age and partly to the development which the animal undergoes during the progress of her initial official test. This latter, which has been called the factor of development, averages 11.0 per cent of the reentry record for Jerseys and 12.2 per cent for Guernseys.

The average initial records of cows retested are somewhat above the average for all cows having initial records.

In the case of the Jersey records studied, carrying a calf for 155 days of the test period has no noticeable effect on the average production of heifers up to the age of 3 years, but the average production of cows 3 years old and upward which met this requirement was less than that of the cows which failed to qualify.

BUREAU OF HOME ECONOMICS
RECEIVED
AUG 31 1925
OFFICE OF THE CHIEF

ADDITIONAL COPIES
OF THIS PUBLICATION MAY BE PROCURED FROM
THE SUPERINTENDENT OF DOCUMENTS
GOVERNMENT PRINTING OFFICE
WASHINGTON, D. C.
AT
5 CENTS PER COPY